

UNIT TERMINAL OBJECTIVE

6-1 At the completion of this unit, the EMT-Intermediate student will be able to apply utilize the assessment findings to formulate and implement a treatment plan for a normal or abnormal labor.

COGNITIVE OBJECTIVES

At the completion of this unit, the EMT-Intermediate student will be able to:

- 6-1.1 Review the anatomic structures and physiology of the reproductive system. (C-1)
- 6-1.2 Identify the normal events of pregnancy. (C-1)
- 6-1.3 Describe how to assess an obstetrical patient. (C-1)
- 6-1.4 Identify the stages of labor and the EMT-Intermediate's role in each stage. (C-1)
- 6-1.5 Differentiate between normal and abnormal delivery. (C-3)
- 6-1.6 Identify and describe complications associated with pregnancy and delivery. (C-1)
- 6-1.7 Identify predelivery emergencies. (C-1)
- 6-1.8 State indications of an imminent delivery. (C-1)
- 6-1.9 Differentiate the management of a patient with predelivery emergencies from a normal delivery. (C-3)
- 6-1.10 State the steps in the predelivery preparation of the mother. (C-1)
- 6-1.11 State the steps to assist in the delivery of a newborn. (C-1)
- 6-1.12 Describe how to care for the newborn. (C-1)
- 6-1.13 Describe how and when to cut the umbilical cord. (C-1)
- 6-1.14 Discuss the steps in the delivery of the placenta. (C-1)
- 6-1.15 Describe the management of the mother post-delivery. (C-1)
- 6-1.16 Describe the procedures for handling abnormal deliveries. (C-1)
- 6-1.17 Describe the procedures for handling complications of pregnancy. (C-1)
- 6-1.18 Describe the procedures for handling maternal complications of labor. (C-1)
- 6-1.19 Describe special considerations when meconium is present in amniotic fluid or during delivery. (C-1)
- 6-1.20 Describe special considerations of a premature baby. (C-1)

AFFECTIVE OBJECTIVES

At the completion of this unit, the EMT-Intermediate student will be able to:

- 6-1.21 Advocate the need for treating two patients (mother and baby). (A-2)
- 6-1.22 Value the importance of maintaining a patient's modesty and privacy during assessment and management. (A-2)
- 6-1.23 Serve as a role model for other EMS providers when discussing or performing the steps of childbirth. (A-3)
- 6-1.24 Value the importance of body substance insolation. (A-2)

PSYCHOMOTOR OBJECTIVES

At the completion of this unit, the EMT-Intermediate student will be able to:

- 6-1.25 Demonstrate how to assess an obstetric patient. (P-2)
- 6-1.26 Demonstrate how to provide care for a patient with: (P-2)
 - a. Excessive vaginal bleeding
 - b. Abdominal pain
- 6-1.27 Demonstrate how to prepare the obstetric patient for delivery. (P-2)
- 6-1.28 Demonstrate how to assist in the normal cephalic delivery of the fetus. (P-2)
- 6-1.29 Demonstrate how to deliver the placenta. (P-2)
- 6-1.30 Demonstrate how to provide post-delivery care of the mother. (P-2)
- 6-1.31 Demonstrate how to assist with abnormal deliveries. (P-2)

- a. Membrane surrounding fetus
 - b. Fluid originates from fetal sources - urine, secretions
 - c. Between 500 and 1000 cc's of fluid after 20 weeks
 - d. Rupture of the membrane produces watery discharge
- C. Fetal growth process
- 1. End of 3rd month
 - a. Sex may be distinguished
 - b. Heart is beating
 - c. Every structure found at birth is present
 - 2. End of 5th month
 - a. Fetal heart tones can be detected
 - b. Fetal movement may be felt by the mother
 - 3. End of 6th month
 - a. May be capable of survival if born prematurely
 - 4. Approximately middle of 10th month
 - a. Considered to have reached full term
 - b. Expected date of confinement (EDC)
- D. Obstetric terminology
- 1. Antepartum - before delivery
 - 2. Postpartum - after delivery
 - 3. Prenatal - existing or occurring before birth
 - 4. Natal - connected with birth
 - 5. Gravida - number of pregnancies
 - 6. Para - number of pregnancies carried to full term
 - 7. Primigravida - a woman who is pregnant for the first time
 - 8. Primipara - a woman who has given birth to her first child
 - 9. Multiparous - a woman who has given birth multiple times
 - 10. Gestation - period of time for intrauterine fetal development

III. General assessment of the obstetric patient

- A. Initial assessment
- B. History of present illness
 - 1. SAMPLE
 - a. Pertinent medical history
 - (1) Diabetes
 - (2) Heart disease
 - (3) Hypertension/ hypotension
 - (4) Seizures
 - 2. Current health of patient
 - a. Pre-existing conditions
 - b. Prenatal care
 - (1) None
 - (2) Physician
 - (3) Nurse midwife
 - c. Illicit drug use
- C. Obstetrical history
 - 1. Length of gestation
 - 2. Primipara or multiparous
 - 3. Previous cesarean sections
 - 4. Previous gynecologic or obstetric complications

- (a) Evaluate impending shock - check orthostatic vital signs
 - (b) Presence and volume of vaginal blood
 - (c) Presence of tissue or large clots
 - d. Transport considerations
 - (1) Collect and transport any passed tissue, if possible
 - e. Psychological support/ communications strategies
 - (1) Emotional support extremely important
- 2. Ectopic pregnancy
 - a. Incidence
 - (1) Approximately 1 of every 200 pregnancies
 - (2) Most are symptomatic and/ or detected 2-12 weeks gestation
 - b. Cause
 - (1) Ovum develops outside the uterus
 - (a) Previous surgical adhesions
 - (b) Pelvic inflammatory disease
 - (c) Tubal ligation
 - (d) Use of an IUD
 - c. Organs affected
 - (1) Fallopian tube
 - d. Complications
 - (1) May be life-threatening
 - (2) May lead to hypovolemic shock and death
 - e. Specific assessment findings
 - (1) Severe abdominal pain, may radiate to back
 - (2) Amenorrhea - absence of monthly blood flow and discharge
 - (3) Vaginal bleeding absent or minimal
 - (4) Upon rupture, bleeding may be excessive
 - (5) Shock signs and symptoms
 - (6) Additional history
 - (a) Previous surgical adhesions
 - (b) Pelvic inflammatory disease
 - (c) Tubal ligation
 - (d) Use of an IUD
 - (e) Previous ectopic pregnancy
 - (7) Additional physical examination
 - (a) Check for impending shock - orthostatic vital signs
 - (b) Presence and volume of vaginal blood
 - f. Additional management
 - (1) See "general management"
 - (2) Second large bore IV line
 - (3) Trendelenburg, if shock impending
 - g. Transport considerations
 - (1) Emergency transport to nearest surgically-capable facility
 - h. Psychological support/ communications strategies
- 3. Placenta previa
 - a. Incidence
 - (1) About 1 in 300
 - (2) Higher in preterm births
 - b. Cause
 - (1) Placenta implantation in lower uterus; covering cervix opening

- d. Cardiac disorders
 - (1) Additional stress on the heart
 - (a) Cardiac output increases 30% by week 34
- 2. Medical complications of pregnancy
 - a. Toxemia (pre-eclampsia/ eclampsia)
 - (1) Incidence
 - (a) Serious condition
 - (b) Pregnancy induced hypertension (PIH)
 - (2) Cause
 - (a) Associated with first birth, multiple births, excessive amniotic fluid
 - (b) Pre-existing conditions
 - i) Hypertension
 - ii) Renal disease
 - iii) Diabetes
 - (3) Organs affected
 - (4) Complications
 - (a) Convulsions seriously threaten the fetus by abruptio placenta
 - (5) Specific assessment findings
 - (a) Occurs in the last trimester of pregnancy
 - (b) Pre-eclampsia is non-convulsive state of toxemia
 - (c) Pre-eclampsia has two of the following three signs
 - i) Hypertension (BP > 140/90 - acute systolic rise > 20 and diastolic rise > 10)
 - ii) Fluid retention with excessive weight gain
 - iii) Proteinuria
 - (d) Eclampsia includes convulsions
 - (e) Additional history
 - i) Hypertension
 - ii) Excessive weight gain with edema and/ or seizures
 - (f) Additional physical exam
 - i) Headaches and/ or epigastric pain; possible seizure
 - ii) Visual problems
 - (6) Transport considerations
 - (a) If a seizure has not occurred
 - i) Keep patient calm and quiet
 - ii) IV access
 - iii) Darken ambulance
 - iv) Position patient LLR
 - v) Transport gently
 - vi) Minimize stimuli to avoid precipitating seizure
 - (b) If a seizure is occurring
 - i) IV access
 - ii) Consider the administration of 5 to 10 mg of diazepam IV push
 - (c) Emergency transport to appropriate facility
 - (d) Definitive treatment is cesarean section
 - (7) Psychological support/ communications strategies
 - b. Diabetes

- a. Related to the imminence of delivery
 - (1) Number of pregnancies
 - (a) Labor is shortened with multiparity
 - (2) Frequency of contractions
 - (a) Two minutes apart may signal imminent delivery
 - (3) Maternal urge to push
 - (a) Desire to push signals imminent delivery
 - (4) Crowning of the presenting part
 - (a) Imminent delivery
 - b. Related to the presence of complications
 - (1) Abnormal presentation
 - (2) Fetal distress
 - (3) Multiple births
2. Delivery of the newborn
 - a. Prepare a delivery area
 - (1) Clean, adequate space
 - b. Provide oxygen to the mother
 - (1) Nonrebreather or nasal cannula
 - c. Establish an IV
 - (1) KVO/ TKO rate
 - d. Position mother on her back and drape appropriately
 - e. Monitor the fetal heart rate, if time allows
 - f. Coach the mother in breathing patterns
 - g. Encourage mother to push with contractions
 - h. Establish body substance isolation
 - i. Control the delivery of the fetal head
 - (1) Apply gentle hand pressure on the head
 - (2) Beware of fontanelle
 - (3) Support the head as it delivers
 - j. Tear amniotic sac if it continues to cover the baby's head
 - (1) Permits escape of amniotic fluid
 - (2) Allows the newborn to start breathing
 - k. Check for the presence of the umbilical cord wrapped around the neck
 - (1) Carefully remove it
 - l. Suction the neonate's mouth and nose
 - m. Provide support as the head rotates and the shoulders deliver
 - (1) Keep the neonate's head above the level of the vagina
 - n. Clamp the umbilical cord
 - (1) First clamp approximately 4 inches from the neonate
 - (2) Second clamp approximately 6 inches from the neonate
 - (3) Cut the cord between the two clamps
 - o. Support and evaluate the neonate following delivery
 3. Delivery of the placenta
 - a. Usually occurs 5-20 minutes after delivery of neonate
 - b. Do not delay transport to wait for the delivery of the placenta
 - c. If it delivers, place the placenta in a plastic bag

E. Additional care

1. Care for the mother
 - a. Excessive bleeding
 - (1) Perform fundal massage of the uterus

- a. Loss of more than 500 cc's of blood immediately following delivery
- b. May be caused by
 - (1) Lack of uterine tone
 - (2) Vaginal or cervical tears
 - (3) Retained pieces of the placenta
 - (4) Clotting disorders
- c. Incidence
- d. Assessment
 - (1) History
 - (a) Large infant
 - (b) Multiple births have occurred
 - (c) The patient has had placenta previa
 - (d) The patient has had abruptio placenta
 - (e) The patient has had prolonged labor
 - (2) Physical examination
 - (a) Treat the patient, EMT-Intermediate must rely on the patient's clinical appearance and vital signs
 - (b) The uterus feels soft on palpation
 - (c) Inspect the external genitalia for injury resulting in excessive bleeding
 - (d) Observe for signs and symptoms of hypovolemic shock
- e. Management
 - (1) Airway and ventilatory support
 - (a) High flow, high concentration oxygen
 - (2) Circulatory support
 - (3) Pharmacologic interventions
 - (a) Consider 2 large-bore IV's for volume replacement
 - (4) Non-pharmacologic interventions
 - (a) Place the infant at the mother's breast if just delivered
 - (b) Provide uterine massage
 - (c) Do not attempt to force delivery of the placenta
 - (d) Do not pack the vagina
 - (5) Transport considerations
 - (a) Emergent transport of the patient
 - (6) Psychological support/ communications strategies